


Analysis 3 Exercise 8

David Lang
18.11.2022



Outline

1. Serie 7 Review
2. Course Overview
3. Homogeneous Heat Equation
4. Homogeneous Wave Equation
5. Tips for Serie 8

Outline

1. Serie 7 Review

2. Course Overview

3. Homogeneous Heat Equation

4. Homogeneous Wave Equation

5. Tips for Serie 8

Serie 7 Review

1. (Non)homogeneous wave equation
 - Stick to the definition given in the problem.

–

2. Propagation of symmetries from initial data, II

- $u_{tt} - 2u_{xx} = 0$ implies $c = \sqrt{2}$

3. Wave equation on a ring

–

4. Multiple choice

$$g(x) = \sum_{n=0}^N b_n \cos(nx) \text{ and } \int_0^{2\pi} g(x) dx = 0, \text{ then } b_0 = \frac{1}{2\pi} \int_0^{2\pi} g(x) dx = 0$$

.

5. Strange wave equation

–

Outline

1. Serie 7 Review

2. Course Overview

3. Homogeneous Heat Equation

4. Homogeneous Wave Equation

5. Tips for Serie 8

Course Overview

- 1st order PDEs
 - Quasilinear first order PDEs
 - ▶ Method of characteristics
 - ▶ Conservation laws
- 2nd order PDEs
 - Hyperbolic PDEs
 - ▶ Wave equation
 - ▶ D'Alembert formula
 - ▶ **Separation of variables**
 - Parabolic PDEs
 - ▶ **Heat equation**
 - ▶ Maximum principle
 - ▶ **Separation of variables**
 - Elliptic PDEs
 - ▶ Laplace equation
 - ▶ Maximum principle
 - ▶ Separation of variables

Outline

1. Serie 7 Review

2. Course Overview

3. Homogeneous Heat Equation

4. Homogeneous Wave Equation

5. Tips for Serie 8

Homogeneous Heat equation with Dirichlet boundary conditions

Derived in lecture

$$\begin{cases} u_t - ku_{xx} = 0 & (x, t) \in (0, L) \times (0, \infty) \\ u(0, t) = u(L, t) = 0 & t > 0 \\ u(x, 0) = f(x) & x \in (0, L) \end{cases}$$

$$u(x, t) = X(x)T(t)$$

$$u(x, t) = \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi}{L}x\right) e^{-k\left(\frac{n\pi}{L}\right)^2 t}$$

$$f(x) = u(x, 0) = \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi}{L}x\right)$$

$$A_m = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{m\pi}{L}x\right) dx$$

Homogeneous Heat equation with Neumann boundary conditions

$$\begin{cases} u_t - ku_{xx} = 0 & (x, t) \in (0, L) \times (0, \infty) \\ u_x(0, t) = u_x(L, t) = 0 & t > 0 \\ u(x, 0) = f(x) & x \in (0, L) \end{cases}$$

$$u(x, t) = X(x)T(t)$$

$$T'(t)X(x) - kX''(x)T(t) = 0$$

$$\frac{T'(t)}{kT(t)} = \frac{X''(x)}{X(x)} = -\lambda$$

$$X''(x) = -\lambda X(x) \quad \text{and} \quad T'(t) = -\lambda kT(t)$$

$$X(x) = \begin{cases} \alpha \cosh(\sqrt{-\lambda}x) + \beta \sinh(\sqrt{-\lambda}x), & \lambda < 0 \\ \alpha + \beta x, & \lambda = 0 \\ \alpha \cos(\sqrt{\lambda}x) + \beta \sin(\sqrt{\lambda}x), & \lambda > 0 \end{cases}$$

Homogeneous Heat equation with Neumann boundary conditions

$$X'(x) = \begin{cases} \sqrt{-\lambda} [\alpha \sinh(\sqrt{-\lambda}x) + \beta \cosh(\sqrt{-\lambda}x)] , & \lambda < 0 \\ \beta, & \lambda = 0 \\ \sqrt{\lambda} [-\alpha \sin(\sqrt{\lambda}x) + \beta \cos(\sqrt{\lambda}x)] , & \lambda > 0 \end{cases}$$

If $\lambda < 0$, then $X(x) = 0$

If $\lambda = 0$, then $X(x) = \alpha$

If $\lambda > 0$, then $X_n(x) = \alpha_n \cos(\sqrt{\lambda_n}x)$ and $\lambda_n = (\frac{n\pi}{L})^2$

$$T'(t) = -\lambda k T(t)$$

$$T_n(t) = B_n e^{-k(\frac{n\pi}{L})^2 t}$$

$$u(x, t) = \frac{1}{2}B_0 + \sum_{n=1}^{\infty} B_n \cos(\frac{n\pi}{L}x) e^{-k(\frac{n\pi}{L})^2 t}$$

$$f(x) = u(x, 0) = \frac{1}{2}B_0 + \sum_{n=1}^{\infty} B_n \cos(\frac{n\pi}{L}x)$$

$$B_m = \frac{2}{L} \int_0^L f(x) \cos(\frac{m\pi}{L}x) dx$$

Example 1

$$\begin{cases} u_t - ku_{xx} = 0 & (x, t) \in (0, \pi/2) \times (0, \infty) \\ u(0, t) = u(\pi/2, t) = 0 & t > 0 \\ u(x, 0) = 3 \sin(4x) & x \in (0, L) \end{cases}$$

Outline

1. Serie 7 Review
2. Course Overview
3. Homogeneous Heat Equation
- 4. Homogeneous Wave Equation**
5. Tips for Serie 8

Homogeneous Wave equation with Neumann boundary conditions

Derived in Lecture

$$\begin{cases} u_{tt} - c^2 u_{xx} = 0 & (x, t) \in (0, L) \times (0, \infty) \\ u_x(0, t) = u_x(L, t) = 0 & t > 0 \\ u(x, 0) = f(x) & x \in (0, L) \\ u_t(x, 0) = g(x) & x \in (0, L) \end{cases}$$

$$u(x, t) = \frac{A_0 + B_0 t}{2} + \sum_{n=1}^{\infty} \cos\left(\frac{n\pi}{L}x\right) \left[A_n \cos\left(\frac{n\pi c}{L}t\right) + B_n \sin\left(\frac{n\pi c}{L}t\right) \right]$$

$$f(x) = u(x, 0) = \frac{A_0}{2} + \sum_{n=1}^{\infty} A_n \cos\left(\frac{n\pi}{L}x\right)$$

$$A_m = \frac{2}{L} \int_0^L f(x) \cos\left(\frac{m\pi}{L}x\right) dx$$

$$g(x) = u_t(x, 0) = \frac{B_0}{2} + \sum_{n=1}^{\infty} B_n \frac{n\pi c}{L} \cos\left(\frac{n\pi}{L}x\right)$$

$$B_0 = \frac{2}{L} \int_0^L g(x) dx, \quad B_m = \frac{2}{cm\pi} \int_0^L g(x) \cos\left(\frac{m\pi}{L}x\right) dx$$

Example 2

$$\begin{cases} u_{tt} - 4u_{xx} = 0 & (x, t) \in (0, 1) \times (0, \infty) \\ u_x(0, t) = u_x(1, t) = 0 & t > 0 \\ u(x, 0) = \cos^2(\pi x) & x \in (0, 1) \\ u_t(x, 0) = \sin^2(\pi x) \cos(\pi x) & x \in (0, 1) \end{cases}$$

Example 2

$$\begin{cases} u_{tt} - 4u_{xx} = 0 & (x, t) \in (0, 1) \times (0, \infty) \\ u_x(0, t) = u_x(1, t) = 0 & t > 0 \\ u(x, 0) = \cos^2(\pi x) & x \in (0, 1) \\ u_t(x, 0) = \sin^2(\pi x) \cos(\pi x) & x \in (0, 1) \end{cases}$$

Homogeneous Wave equation with Dirichlet boundary conditions

$$\begin{cases} u_{tt} - c^2 u_{xx} = 0 & (x, t) \in (0, L) \times (0, \infty) \\ u(0, t) = u(L, t) = 0 & t > 0 \\ u(x, 0) = f(x) & x \in \mathbb{R} \\ u_t(x, 0) = g(x) & x \in \mathbb{R} \end{cases}$$

$$u(x, t) = \sum_{n=1}^{\infty} \sin\left(\frac{n\pi}{L}x\right) \left[A_n \cos\left(\frac{n\pi c}{L}t\right) + B_n \sin\left(\frac{n\pi c}{L}t\right) \right]$$

$$f(x) = u(x, 0) = \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi}{L}x\right)$$

$$A_m = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{m\pi}{L}x\right) dx$$

$$g(x) = u_t(x, 0) = \sum_{n=1}^{\infty} \frac{n\pi c}{L} B_n \sin\left(\frac{n\pi}{L}x\right)$$

$$B_m = \frac{2}{cm\pi} \int_0^L g(x) \sin\left(\frac{m\pi}{L}x\right) dx$$

Example 3

$$\begin{cases} u_{tt} - 4u_{xx} = 0 & (x, t) \in (0, \pi) \times (0, \infty) \\ u(x, 0) = 3 \cos(x) & x \in \mathbb{R} \\ u_t(x, 0) = 1 - \cos(4x) & x \in \mathbb{R} \\ u_x(0, t) = u_x(\pi, t) = 0 & t > 0 \end{cases}$$

Example 3

$$\begin{cases} u_{tt} - 4u_{xx} = 0 & (x, t) \in (0, \pi) \times (0, \infty) \\ u(x, 0) = 3 \cos(x) & x \in \mathbb{R} \\ u_t(x, 0) = 1 - \cos(4x) & x \in \mathbb{R} \\ u_x(0, t) = u_x(\pi, t) = 0 & t > 0 \end{cases}$$

Example 3

$$\begin{cases} u_{tt} - 4u_{xx} = 0 & (x, t) \in (0, \pi) \times (0, \infty) \\ u(x, 0) = 3 \cos(x) & x \in \mathbb{R} \\ u_t(x, 0) = 1 - \cos(4x) & x \in \mathbb{R} \\ u_x(0, t) = u_x(\pi, t) = 0 & t > 0 \end{cases}$$

Outline

1. Serie 7 Review
2. Course Overview
3. Homogeneous Heat Equation
4. Homogeneous Wave Equation
5. Tips for Serie 8

Tips for Serie 8

1. Separation of variables

- I suggest repeating the derivation at least once. Consult the general solution only if you are fully familiar with the procedure.
- (a)
- (b) $4\sin^3(x) = 3\sin(x) - \sin(3x)$
- (c)

2. Multiple choice

- (a) Fourier Series and Orthogonality
- (b) Solve $u(x, t)$ explicitly and explore the trigonometric terms.

More on Waves:

1. AT&T, Dr. Shive: <https://youtu.be/DovunOxIY1k>

More on Fourier Series:

1. Prof. Strang: <https://youtu.be/vA9dfINW4Rg>

References:

1. Lecture notes on the course website.
2. "An Introduction to Partial Differential Equations" by Yehuda Pinchover and Jacob Rubinstein