

4.1. Conservation laws and critical times Consider the PDE

$$u_y + \partial_x(f(u)) = 0.$$

In the following cases, compute the critical time y_c (i.e., the first time when the solution becomes nonsmooth):

(a) $f(u) = \frac{1}{2}u^2$, the initial datum is $u(x, 0) = \sin(x)$.

(b) $f(u) = \sin(u)$, the initial datum is $u(x, 0) = x^2$.

(c) $f(u) = e^u$, the initial datum is $u(x, 0) = x^3$.

4.2. Multiple choice Cross the correct answer(s).

(a) In all generality, a conservation law (as we defined it in the lecture)

- | | |
|---|---|
| <input type="radio"/> admits a strong local solution | <input type="radio"/> has finite critical time |
| <input type="radio"/> admits a strong global solution | <input type="radio"/> might have several weak solutions |
| <input type="radio"/> develops singularities | <input type="radio"/> has straight lines as characteristics |

(b) Consider the conservation law

$$\begin{cases} u_y + (\alpha u^2 - u)u_x = 0, & (x, y) \in \mathbb{R} \times (0, +\infty), \\ u(x, 0) = 1, & x < 0, \\ u(x, 0) = 0, & x \geq 0. \end{cases}$$

Then, the shock wave solution has strictly positive slope if

- | | |
|------------------------------------|--|
| <input type="radio"/> $\alpha > 1$ | <input type="radio"/> $\alpha < 0$ |
| <input type="radio"/> $\alpha < 1$ | <input type="radio"/> $\alpha > \frac{3}{2}$ |
| <input type="radio"/> $\alpha > 0$ | <input type="radio"/> $\alpha < \frac{3}{2}$ |

(c) The conservation law

$$\begin{cases} u_y + (u^2 + 5)u_x = 0, & (x, y) \in \mathbb{R} \times (0, +\infty), \\ u(x, 0) = 1, & x < 0, \\ u(x, 0) = \sqrt{1-x}, & x \in [0, 1], \\ u(x, 0) = 0, & x > 1, \end{cases} \tag{1}$$

has a crossing of characteristics at¹

¹You can partially check your answer computing y_c . Why?

